

Children with spastic Cerebral Palsy experience difficulties adjusting their gait pattern to weight added to the waist, while typically developing children do not

Pieter Meyns^{1*}, Leen Van Gestel², Lynn Bar-on², Marije Goudriaan², Hans Wambacq³, Erwin Aertbeliën³, Herman Bruyninckx³, Guy Molenaers⁴, Paul De Cock⁵, Els Ortibus⁵, Kaat Desloovere⁶

¹Rehabilitation medicine, VUmc, Netherlands, ²Rehabilitation Sciences, KU Leuven, Belgium, ³Mechanical Engineering, KU Leuven, Belgium, ⁴Paediatric Orthopaedics, UZ Leuven, Belgium, ⁵Centre for developmental Disabilities, UZ Leuven, Belgium, ⁶Clinical Motion Analysis Laboratory, UZ Leuven, Belgium

Submitted to Journal:
Frontiers in Human Neuroscience

Article type:
Original Research Article

Manuscript ID:
208469

Received on:
04 May 2016

Revised on:
05 Dec 2016

Frontiers website link:
www.frontiersin.org

Conflict of interest statement

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest

Author contribution statement

LVG and KD conceived and designed the experiment. GM, PDC, EO and LVG performed patient recruitment. PM and LVG tested the participants. PM, LVG and LB performed statistical analyses and interpreted the data. HW, EA, and HB created the software for the data extraction of the three dimensional gait analyses. MG, HW and EA performed data quality checks. PM, LVG, and KD wrote the paper. MG performed the lay-out and edited the paper. The writing process and data analysis were supervised by KD.

Keywords

Cerebral Palsy, Gait, weight, Body mass, EMG, Muscle Weakness

Abstract

Word count: 350

The prevalence of childhood overweight and obesity is increasing in the last decades, also in children with Cerebral Palsy (CP). Even though it has been established that an increase in weight can have important negative effects on gait in healthy adults and children, it has not been investigated what the effect is of an increase in body weight on the characteristics of gait in children with CP. In CP, pre and post three-dimensional gait analyses are performed to assess the effectiveness of an intervention. As a considerable amount of time can elapse between these measurements, and the effect of an alteration in the body weight is not taken into consideration, this effect of increased body weight is of specific importance.

Thirty children with the predominantly spastic type of CP and fifteen typically developing (TD) children were enrolled (age 3 to 15 years). All children underwent three-dimensional gait analysis with weight-free (baseline) and weighted (10% of the body weight added around their waist) trials.

Numerous gait parameters showed a different response to the added weight for TD and CP children. TD children increased walking velocity, step- and stride length, and decreased double support duration with a slightly earlier timing of foot-off, while the opposite was found in CP. Similarly, increased ranges of motion at the pelvis (coronal plane) and hip (all planes), higher joint angular velocities at the hip and ankle, as well as increased moments and powers at the hip, knee and ankle were observed for TD children, while CP children did not change or even showed decreases in the respective measures in response to walking with added weight. Further, while TD children increased their gastrocnemius EMG amplitude during weighted walking, CP children slightly decreased their gastrocnemius EMG amplitude.

As such, an increase in weight has a significant effect on the gait pattern in CP children. Clinical gait analysts should therefore take into account the negative effects of increased weight during pre-post measurements to avoid misinterpretation of treatment results. Overweight and obesity in CP should be counteracted or prevented as the increased weight has detrimental effects on the gait pattern.

Funding statement

Pieter Meyns was supported by a grant of the Special Research Fund of the KU Leuven (OT/08/034 and PMK/12/180) and by FWO (Grant G.0901.11, 'Krediet aan Navorsers' Grant 1503915N). Pieter Meyns is supported by the European Commission (Horizon 2020) as a Marie Skłodowska-Curie fellow (proposal 660458). Leen Van Gestel received a PhD fellowship of the Research Foundation-Flanders (FWO), Belgium. The authors gratefully acknowledge the IPSA project (060799) which was supported by the IWT Flanders. Lynn Bar-On was supported by a Grant from the Doctoral Scholarships Committee for International Collaboration with non EER-countries (DBOF/12/058) of the University of Leuven, Belgium. Marije Goudriaan was supported by the MD-Paedigree project: a Model-Driven Paediatric European Digital Repository, partially funded by the European Commission under P7-ICT-2011-9 programme (600932). Kaat Desloovere is supported for this research by the Research Fund KU Leuven (OT/12/100): "Pattern recognition for gait analysis: integration of clinical expert knowledge and machine learning techniques." There was no role of these funding bodies in the study design, collection and analysis of data, interpretation of data, writing of the paper and decision to submit the paper for publication.

Ethics statements

(Authors are required to state the ethical considerations of their study in the manuscript, including for cases where the study was exempt from ethical approval procedures)

Does the study presented in the manuscript involve human or animal subjects: Yes

Please provide the complete ethics statement for your manuscript. Note that the statement will be directly added to the manuscript file for peer-review, and should include the following information:

- Full name of the ethics committee that approved the study
- Consent procedure used for human participants or for animal owners
- Any additional considerations of the study in cases where vulnerable populations were involved, for example minors, persons with disabilities or endangered animal species

As per the Frontiers authors guidelines, you are required to use the following format for statements involving human subjects: This study was carried out in accordance with the recommendations of 'name of guidelines, name of committee' with written informed consent from all subjects. All subjects gave written informed consent in accordance with the Declaration of Helsinki. The protocol was approved by the 'name of committee'.

For statements involving animal subjects, please use:

This study was carried out in accordance with the recommendations of 'name of guidelines, name of committee'. The protocol was approved by the 'name of committee'.

If the study was exempt from one or more of the above requirements, please provide a statement with the reason for the exemption(s).

Ensure that your statement is phrased in a complete way, with clear and concise sentences.

Medical Ethics Committee UZ KU Leuven / Research

In accordance with the Declaration of Helsinki, written informed consent was obtained of the participants' parents.

The participants parents provided written informed consent prior to the experiment. The participants parents supervised the measurement session.

**Children with spastic Cerebral Palsy experience difficulties
adjusting their gait pattern to weight added to the waist, while
typically developing children do not**

Running title; Weighted walking in CP

Pieter Meyns^{*1*}, Leen Van Gestel^{2*}, Lynn Bar-On^{2,3}, Marije Goudriaan^{2,3}, Hans Wambacq⁴,
Erwin Aertbeliën⁴, Herman Bruyninckx⁴, Guy Molenaers^{3,5,6}, Paul De Cock⁷, Els Ortibus⁷, Kaat
Desloovere^{2,3}

1. Department of Rehabilitation Medicine, MOVE Research Institute Amsterdam, VU
University Medical Center, Amsterdam, The Netherlands

2. KU Leuven, Faculty of Kinesiology and Rehabilitation Sciences, Department of
Rehabilitation Sciences, Belgium

3. Clinical Motion Analysis Laboratory, CERM, University Hospital Leuven, Belgium

4. KU Leuven, Faculty of Engineering, Department of Mechanical Engineering, Division of
Production Engineering, Machine Design and Automation, Belgium

5. Department of Paediatric Orthopaedics, University Hospital Leuven, Belgium

6. KU Leuven, Faculty of Medicine, Department of Musculoskeletal Sciences, Belgium

7. Centre for developmental Disabilities, University Hospital Leuven, Belgium

* Pieter Meyns and Leen Van Gestel are joint-first authors, both authors contributed equally

To be submitted in Frontiers in Human Neuroscience

Research topic; Motor control of gait and the underlying neural network in pediatric
neurology

Type of Article: Original research

*Correspondence:

Pieter Meyns
Department of Rehabilitation Medicine, MOVE Research Institute Amsterdam
VU University Medical Center
De Boelelaan 1118, 1081 HV Amsterdam, The Netherlands
pieter.meyns@gmail.com

Word Count Abstract:	350 (max 350)
Word Count Main Text:	3945 (max 12000)
Number of Tables and Figures:	6 (max 15)

Number of Appendices: 1

Author contribution statement

LVG and KD conceived and designed the experiment. GM, PDC, EO and LVG performed patient recruitment. PM and LVG tested the participants. PM, LVG and LB performed statistical analyses and interpreted the data. HW, EA, and HB created the software for the data extraction of the three dimensional gait analyses. LVG, MG, HW and EA performed data quality checks. PM, LVG, and KD wrote the paper. MG performed the lay-out and edited the paper. The writing process and data analysis were supervised by KD.

Funding

Pieter Meyns was supported by a grant of the Special Research Fund of the KU Leuven (OT/08/034 and PMK/12/180) and by FWO (Grant G.0901.11, ‘Krediet aan Navorsers’ Grant 1503915N). Pieter Meyns is supported by the European Commission (Horizon 2020) as a Marie Skłodowska-Curie fellow (proposal 660458). Leen Van Gestel received a PhD fellowship of the Research Foundation-Flanders (FWO), Belgium. The authors gratefully acknowledge the IPSA project (060799) which was supported by the IWT Flanders. Lynn Bar-On was supported by a Grant from the Doctoral Scholarships Committee for International Collaboration with non EER-countries (DBOF/12/058) of the University of Leuven, Belgium. Marije Goudriaan was supported by the MD-Paedigree project: a Model-Driven Paediatric European Digital Repository, partially funded by the European Commission under P7-ICT-2011-9 programme (600932). Kaat Desloovere is supported for this research by the Research Fund KU Leuven (OT/12/100): “Pattern recognition for gait analysis: integration of clinical expert knowledge and machine learning techniques.” There was no role of these funding bodies in the study design, collection and analysis of data, interpretation of data, writing of the paper and decision to submit the paper for publication.

Acknowledgements

The authors would like to thank Steffen Fieuws of the Leuven Biostatistics and Statistical Bioinformatics Centre for his expertise and advice on the statistical analyses of this paper.

Abstract

The prevalence of childhood overweight and obesity is increasing in the last decades, also in children with Cerebral Palsy (CP). Even though it has been established that an increase in weight can have important negative effects on gait in healthy adults and children, it has not been investigated what the effect is of an increase in body weight on the characteristics of gait in children with CP. In CP, pre and post three-dimensional gait analyses are performed to assess the effectiveness of an intervention. As a considerable amount of time can elapse between these measurements, and the effect of an alteration in the body weight is not taken into consideration, this effect of increased body weight is of specific importance.

Thirty children with the predominantly spastic type of CP and fifteen typically developing (TD) children were enrolled (age 3 to 15 years). All children underwent three-dimensional gait analysis with weight-free (baseline) and weighted (10% of the body weight added around their waist) trials.

Numerous gait parameters showed a different response to the added weight for TD and CP children. TD children increased walking velocity, step- and stride length, and decreased double support duration with a slightly earlier timing of foot-off, while the opposite was found in CP. Similarly, increased ranges of motion at the pelvis (coronal plane) and hip (all planes), higher joint angular velocities at the hip and ankle, as well as increased moments and powers at the hip, knee and ankle were observed for TD children, while CP children did not change or even showed decreases in the respective measures in response to walking with added weight. Further, while TD children increased their gastrocnemius EMG amplitude during weighted walking, CP children slightly decreased their gastrocnemius EMG amplitude.

As such, an increase in weight has a significant effect on the gait pattern in CP children. Clinical gait analysts should therefore take into account the negative effects of increased weight during pre-post measurements to avoid misinterpretation of treatment results. Overweight and obesity in CP should be counteracted or prevented as the increased weight has detrimental effects on the gait pattern.

Keywords

Cerebral Palsy, gait, weight, body mass, EMG, muscle weakness

1. Introduction

According to the World Health Organization (WHO), in 2014 more than 1.9 billion adults were overweight. Of these over 600 million were obese (World Health Organization, 2016b). Furthermore, the worldwide prevalence of obesity has more than doubled between 1980 and 2013 (Finucane et al., 2011; Ng et al., 2014).

Obesity has an important impact on reduced quality of life (Taylor et al., 2013) and public health (Visscher and Seidell, 2001), as it is related to the development of several non-communicable diseases such as cardiovascular disease, type 2 diabetes mellitus, and osteoarthritis (World Health Organization, 2016b).

Similarly, in children an increase in overweight and obesity has been observed over the last decades (de Onis et al., 2010). This is not only the case in typically developing children (TDC), but also in children that already present with a pathology; e.g. children with Cerebral Palsy (CP) (Park et al., 2011; Rogozinski et al., 2007).

Cerebral Palsy is the most common developmental cause of physical disability in the world (Aisen et al., 2011), with a prevalence of 2-3 in 1000 live births (Odding et al., 2006). It originates from non-progressive impairments to the brain before, during or shortly after birth (Bax et al., 2005), resulting in persistent (primary) motor and sensory impairments such as abnormal motor control, muscle strength, and/or muscle tone. As a result of abnormal muscle activity or loading of the bones, secondary impairments can develop over time, such as shortened muscles which will restrict the joint range of motion. These primary and secondary impairments result in a pathological gait pattern in CP.

Many treatment modalities in CP are aimed at improving the gait pattern, by focusing on the primary and secondary impairments. Orthoses, for instance, are often used to control the position and movement of the ankle (Novacheck et al., 2009). Neurosurgery, on the other hand, can reduce the patient's spasticity by means of selective dorsal rhizotomy or intrathecal baclofen, while orthopedic surgery can address bone deformities and muscle contractures (Novacheck and Gage, 2007). Gait analysis has a crucial role in the treatment of gait impairments in CP, as it allows 1) the identification of the specific deficits of each patient, and 2) the evaluation of the outcome of treatment interventions specifically selected to target those deficits (Novacheck and Gage, 2007; Schwartz et al., 2004).

Interventions in CP are often assessed via pre-post treatment three-dimensional gait analysis. Until now though, the effect of an alteration in the body weight is usually not taken into consideration when interpreting pre-post data, even though a considerable amount of time can elapse between the two measurements.

However, literature in other populations provides indications that changes in body weight result in alterations in the gait pattern. In children with overweight and obesity for example, it has been reported that additional body mass leads to higher foot loading, with a disproportional impact on the midfoot area (Cousins et al., 2013; Mueller et al., 2016). Both adults and children with obesity have been found to walk with increased peak knee adduction moments (which may result in excessive joint loading) (Browning and Kram, 2007; Gushue et al., 2005). Adults with obesity also have been reported to walk with increased external knee flexion moments, and decreased knee rotation moments compared to non-overweight adults (Harding et al., 2012).

Furthermore, some researchers have indicated changes in EMG activations due to obesity in children and adults as well. Blakemore et al. (2013), showed that during fast walking, overweight children had longer gastrocnemius activity during stance, but shorter

gastrocnemius activity during swing (Blakemore et al., 2013). A study by Amiri et al. (2015) indicated that adults with obesity walked with significantly prolonged gastrocnemius and quadriceps EMG activity during the stance phase (Amiri et al., 2015). In adults, walking while carrying load resulted in increased EMG activations of the vastus lateralis and gastrocnemius, as well as increased durations of biceps femoris, gastrocnemius, vastus lateralis and semimembranosus activity (Ghori and Luckwill, 1985; Simpson et al., 2011).

As such it appears that increased weight has a significant effect on several aspects of the gait pattern, including kinematics, kinetics and EMG.

To the best of our knowledge, the effect of increasing the body weight on the characteristics of gait in CPc has not yet been investigated. To examine this experimentally, one could either gradually increase the weight of CPc over several weeks, or investigate the effect of an immediate increase in weight. One could argue that the gait adaptations to an immediate increase of weight will not necessarily resemble those that happen during the slow increase in body weight as is the case for obesity. In the latter case, the gait pattern can slowly adapt to very small increments in weight. However, body weight can increase more rapidly during daily life as well i.e. during the growth spurt. During the growth spurt, ambulatory CPc typically show fast increases in body weight, similarly as TDc (Day et al., 2007). In the current study we have opted for the latter, as this is the first step to take to provide further insights into the possible adaptations in CPc to added weight. Thus, the experimental paradigm used in the current study, more closely resembles this phenomenon rather than a slow increase in body weight. As CPc often undergo treatment interventions at an age before or close to the growth spurt, it is of interest to investigate the influence of an immediate increase in weight on the gait parameters in CPc.

The current study examined whether experimentally increasing the body weight influences the spatio-temporal gait characteristics, three-dimensional gait kinematics, kinetics, and EMG activations in CPc. We hypothesize that CPc will experience difficulties in adjusting their gait pattern to the added weights; i.e. they will present with more negative changes in their gait due to the increase of body weight compared to TDc (e.g. decrease walking speed, smaller step lengths and ranges of joint motion). Additionally, we hypothesize that more negative changes in the gait pattern will appear in CPc with relatively weaker lower limb muscles compared to children with relatively strong lower limb muscles as they will experience more difficulties negotiating the increased load imposed on their musculoskeletal system.

2. Materials and Methods

2.1 Patients

For this study, a group of CPc meeting specific inclusion criteria (*CP group*) was compared to a control group of TDc. The characteristics of both groups are discussed below and summarized in Table 1. A schematic overview of the study design can be found in Figure 1.

Table 1 near here

For the *CP group*, CPc were recruited at the Laboratory of Clinical Motion Analysis of the University Hospital Leuven (UZ Pellenberg), when they met the following inclusion criteria: (a) predominantly spastic type of CP, (b) age between 3 and 15 years, (c) ability to walk (with or without walking aids), (d) sufficient cooperation to follow verbal instructions, (e) no history of lower limb surgery, (f) no lower limb Botulinum Toxin-A treatment within 6 months prior to the 3DGA, and (g) adequate or moderate lower limb muscle strength. The latter was defined as a MMT-score of at least >2.5 on a 9-point scale (Table A, Supplementary Data) based on the MMT scale developed by Daniels & Worthingham's scale for manual muscle testing (Cuthbert and Goodheart, 2007; Daniels and Worthingham, 1986). Strength was assessed for the knee flexors, knee extensors, and plantarflexors, as these muscles are assumed to be the main actuators for handling the additional load during the weighted walking (weighted walking task is described below). Based on the median MMT-score for these three muscle groups, the CPc were subdivided into two muscle strength groups: children whose median MMT ≥ 4 were classified into the *adequate* lower limb muscle strength group, while children with a median MMT-score between 2.5 and 4 were classified into the *moderate* lower limb muscle strength group. This cut-off value was selected based on the results of initial pilot tests where it was observed that children with a MMT score > 2.5 were still able to accurately execute the weighted walking task for the full testing procedure. In addition to strength, the selectivity of the three lower limb muscles was assessed using a five-point scale as described by Trost (Trost, 2009) (Table A, Supplementary Data).

Figure 1 near here

Thirty CPc were enrolled in the *CP group*. Their average age was 8y 5m ($\pm 2y$ 3m). Twelve children were classified as unilaterally involved while 18 were bilaterally involved. Seventeen children were classified as GMFCS level I and 13 as level II. Seventeen children had *adequate* lower limb muscle strength, while 13 showed *moderate* lower limb muscle strength. Specific lower limb ranges of motion and levels of selective motor control are summarized in Table 1.

The TD control group consisted of 15 children with an average age of 8y 6m ($\pm 1y$ 3m) without a history of orthopedic or neurological pathology. Comparison of the gait adaptations in response to the weighted walking task in the *CP group* to those observed in the TD group enabled the identification of 'non-typical responses' (gait adaptations that were not observed in the TD group). Non-typical responses unmask 3D gait parameters that alter due to the addition of weight which are specific to the *CP group*.

To avoid the inclusion of correlated data we studied one of the lower limbs in each participant. Therefore, in CPc with unilateral involvement only the involved lower limb side was considered, while in children with bilateral involvement the most involved side was selected. When both sides were equally involved, the left or right lower limb was randomly selected. For TDc, only the right lower limb was investigated.

All experiments were approved by the local ethical committee and informed, written consent was obtained from the study subjects' parents.

2.2 3D gait analysis and weighted walking task

Gait was evaluated through 3D gait analysis (3DGA). Prior to the 3DGA, lower limb dimensions, body height, and weight were measured to enable an estimation of joint center locations and segmental inertia parameters. Patients walked on a 10-meter walkway at a self-selected speed. Spatiotemporal and kinematic measurements were collected using an eight-camera VICON system (612 data capturing system measuring at 100Hz; VICON, Oxford Metrics, Oxford, UK), with the lower limb Plug-In-Gait marker set (Kadaba et al., 1990). Two force plates (Advanced Mechanical Technology Inc., Watertown, MA) were embedded in the walkway for force registration enabling calculation of kinetics. Surface EMG data were collected from the medial hamstrings, rectus femoris, and lateral gastrocnemius muscles using a 16 channel EMG system (K-lab, Biometrics, The Netherlands; or Zerowire, Aurion, Italy), synchronized with the motion capture system. EMG data was collected with a sampling rate of 1500 Hz and filtered with a zero-phase 6th order butterworth filter with a passband ranging from 20 to 500Hz. Nexus software (Oxford Metrics, Oxford, UK) was used to define the gait cycles, to calculate the spatiotemporal parameters, and to estimate the joint angles and internal moments and powers normalized to body mass. Seventy-one discrete parameters were extracted from the continuous kinematic and kinetic waveforms based on a study of the literature (Goldberg et al., 2006; Inman et al., 1981; Perry, 1992; Schutte et al., 2000; Winter, 1987; Wolf et al., 2006) and the routine gait analysis protocol used for children with CP at the Clinical Movement Analysis Laboratory of the University Hospital Leuven (Pellenberg).

All children walked barefoot (baseline/weight-free walking). In addition, all children executed the weighted walking task consisting of barefoot walking with 10% of the body weight added around the waist. This percentage of body weight was selected based on initial pilot tests (n=9). In these tests, it was observed that adding 10% of the body weight resulted in an increased load on the lower limbs without overloading, assuring the ability to walk across the gait lab for the full test procedure. The results from these pilot tests indicated that adding more than 10% of the body weight limited some patients to perform the walking trails. Hence, 10% of the body weight was chosen. The weight was applied evenly around the waist by means of a belt. The belt was positioned close to the level of S2 (approximate location of the centre of gravity) to avoid challenging the equilibrium of the children, and above the marker of the pelvis to ensure full visibility of the lower limb Plug-In-Gait marker set. For both conditions, the children walked at a self-selected speed until three trials with full kinematics, kinetics and EMG were recorded. Internal moments and powers for the weighted walking trials were normalized to the child's body weight plus the additional weight.

2.3 Statistical analysis

For this study, a subset of 71 gait parameters was selected from the output of the 3DGA for both walking conditions (baseline and weighted walking). Spatiotemporal parameters, discrete values of joint angles, moments and powers at specific points in the gait cycle, and the mean EMG frequency over one full gait cycle were automatically extracted from the gait waveforms by means of custom-made software implemented in MATLAB (Mathworks). All parameters were averaged per walking condition over the three registered walking trials. Additionally, the

percentage increase/decrease in mean EMG amplitude (raw signal) in response to the added weight was calculated. All EMG parameters were investigated for the medial hamstrings, rectus femoris, and lateral gastrocnemius muscles.

The difference scores between baseline walking and weighted walking were calculated for all the 3DGA parameters of the *CP group* and the TD group. These scores illustrate how the children responded to an increase in body weight. A Kruskal-Wallis test with post-hoc Mann-Whitney U tests (with False Discovery Rate (FDR) correction for multiple testing) compared the responses of the TD children to the responses of (1) all CPc in the *CP group*, (2) CPc with adequate strength, and (3) CPc with moderate strength. A non-parametric method was applied as not all outcome parameters were characterized by a normal or Gaussian distribution. All statistical procedures were performed with the SAS system (SAS Institute Inc, SAS Campus, Dr. Cary, NC 27513). Level of significance was set at $P < 0.05$.

In review

3. Results

When children walked with added weight, changes in the gait pattern were observed for both TDc and CPc. However, for several 3DGA parameters, CPc responded significantly different to the added weight when compared to TDc. Tables 2 and 3 provide detailed results.

Overall, the TDc appeared to increase their walking velocity, step- and stride length, and decreased their duration of double support while the timing of foot-off decreased slightly, while the opposite pattern was found in CPc (Figure 2 & Table 2).

Figure 2 near here

Increased ranges of motion at the pelvis (coronal plane) and hip (all planes), as well as higher joint angular velocities at the hip and ankle appeared for TDc, while CPc did not seem to change or even decreased the respective ranges of motion and joint angular velocities in response to walking with added weight (Figure 2). Furthermore, several moments and powers at the hip, knee and ankle appeared to increase in TDc, while CPc mainly seemed to decrease their moments and powers during weighted walking (Table 2 & Table 3).

Table 2 near here

Table 3 near here

Finally, a significantly different response in EMG amplitude could be observed in the gastrocnemius. In general, TDc increased their gastrocnemius EMG amplitude when they walked with added weight, while CPc slightly decreased their gastrocnemius EMG amplitude (Figure 3).

Figure 3 near here

When the TD group was compared to the subgroups representing the adequately and moderately strong CP children separately, the response of four 3DGA parameters (timing of foot-off, pelvic range of motion in the coronal plane, maximal hip flexion in swing and maximal hip extension moment in stance) was found to differ significantly confirming the above described results (Tables 2 and 3). No differences in gait parameters due to the increased weight were found between the two CP groups (the adequately and moderately strong CPc).

4. Discussion

Walking with extra weight applied around the waist resulted in multiple changes in the gait pattern of both TDc and CPc. For several 3DGA parameters, these changes were significantly different between the TDc and the *CP group*.

The current results are of clinical importance as, due to the increase in prevalence of overweight and obesity in CPc over the last decades (Park et al., 2011; Rogozinski et al., 2007), the gait pattern in CPc to date can be negatively influenced. Consequently, when interpreting 3D gait analysis data, weight should also be considered as a potential factor influencing the gait parameters.

While TDc seemed to be able to successfully handle the extra weight, CPc experienced difficulties to walk with the increased demand. TDc walked with larger ranges of joint motion, and higher moments and powers in response to the added weight (Table 2, Figure 2). In fact, it appeared that TDc even overcompensated for the extra weight as their resulting gait was faster and more dynamic (reflected in increased kinetics). In contrast, CPc walked slower, with smaller step lengths and ranges of joint motion, and decreased moments and powers. Furthermore, while in TDc the EMG amplitude of the gastrocnemius muscle increased as a result of the added weight, in CPc the EMG amplitude slightly decreased (Figure 3). To increase its force output, a muscle can apply two strategies (or a combination of both). Either more motor units are recruited, or the activity of already recruited motor units is synchronized. The first strategy will increase both the EMG frequency and amplitude, while the latter decreases the EMG frequency whilst increasing the amplitude (Lauer et al., 2008). As the EMG amplitude slightly decreased in CPc, it appears that in CP the muscles failed to successfully increase their force output during weighted walking. This finding is supported by the conclusions in the study of Dallmeijer et al. (2011) who found that the lower limb muscles of CPc are characterized by a small muscle reserve during walking (Dallmeijer et al., 2011).

When the CP group was further subdivided into children with adequate and moderate lower limb muscle strength, no significant differences in 3DGA parameters were found due to the added weight in the two subgroups. As such, it appears that the difference in muscle strength between the two CP strength subgroups did not lead to a significantly different response to the added weight. Nevertheless, both CP subgroups presented with difficulties to walk as a result of an increase in weight.

Previous research already provided an indication that weight could have a negative impact on the gait pattern in CPc. Specifically, the sudden increase in body weight during the pubertal growth spurt has been previously related to a more rapid development of the crouch gait pattern in children with diplegic CP (Gage, 2004). This rapid development was suggested to be related to increased body mass with an unfavorable mass-to-strength ratio; i.e. weak muscles can no longer support the toe walking pattern due to the sudden increase in weight (Kedem and Scher, 2016).

The finding that an increase in body weight has significant negative consequences on several gait parameters in CPc that are frequently used in clinical gait analysis (and the overall gait pattern) is of high clinical importance. Given the fact that interventions in CPc are often evaluated via pre- and post 3D-gait analyses, clinical gait analysts should take into account the increase in weight of the patient, definitely if the assessments take place during or close to the growth spurt. If not, unfavorable (or less favorable than expected) gait outcomes could be related to the treatment rather than to the increase in body weight.

These findings underscore the importance of initiatives such as that of the European Union and the World Health Organization to attempt to cease the rise in overweight and obesity in children (European Union, 2014; World Health Organization, 2016a). These initiatives are of vital importance as they can halt the related health problems associated to overweight, but also stop the negative influence of overweight on the gait pattern in CPc.

When interpreting the results of the current study, one should consider some possible limitations.

It could be argued that similar changes in spatiotemporal parameters of gait have been previously reported as specific markers of balance problems (Liao et al., 1997). However, the weighted walking task of the current study was specifically designed to increase the body weight without challenging the equilibrium by adding the weight evenly around the waist and close to the centre of gravity. Nevertheless, it cannot be ruled out that there was no effect on balance. Further research could focus on further elucidating the individual role of balance on (spatiotemporal) gait parameters during weighted walking.

The decrease in EMG amplitude in CPc could also be partly related to muscle fatigue. It has been described in healthy adults that muscle fatigue influences the EMG signal and e.g. reduces its frequency (Rogers and MacIsaac, 2011). Fatigue could have played a part in the setup of the current study, as the muscle endurance during submaximal tasks in CPc is reduced compared to TDc (Eken et al., 2014). Hence, the combination of the different walking trials with and without weights, and the reduced capacity of CPc to endure activities could have induced muscle fatigue. Combined with the fact that CPc have been shown to have a small muscle reserve during walking (Dallmeijer et al., 2011), this means that the EMG amplitude could not be increased in CPc. Further research examining the mean EMG frequencies and amplitudes of lower limb muscles during gait for several consecutive steps of a large group of CP children might further elucidate this problem.

In the current study, the participants' response was investigated when the weight of the body was experimentally/artificially increased. This allowed us to investigate fundamental changes in the gait pattern in TDc and CPc as a response to a sudden increase in body weight. It could, however, be argued that an increase in body weight in daily life (as happens with overweight and obesity) occurs over a long period of time in which the gait pattern can slowly adapt. Even though the participants in the current study received some practice trials with the weights (until they indicated to feel comfortable with the weights), it is possible that a longer period of adaption could have an impact on the gait pattern changes. Nevertheless, results from previous studies in people with overweight and obesity support our findings (e.g. increase in EMG activations, and knee power generation in TDc). Furthermore, our experimental setup could be specifically interesting in the case of a more rapid increase in body weight which occurs during the growth spurt (Day et al., 2007). In this case, however, one should consider that the current setup does not take into account the combination of the lengthening of the body combined with the increase in body weight. Hence, the weight increase during the growth spurt is spread over the body rather than focused on the trunk.

In *conclusion*, the results of the current paper indicate that a multitude of gait parameters (including spatiotemporal measures, kinematics, kinetics, and EMG) are significantly influenced by adding 10% body weight during walking in TDc and CPc. Furthermore, many of

these gait parameters differed significantly between TDc and CPc in response to the added weight. Given the rapid increase in body weight during the growth spurt in CPc, specialists in clinical gait analysis should take into account the negative effect of the increased body weight during pre-post measurements to avoid misinterpretation of the treatment results. Overweight and obesity in CPc should be counteracted or prevented as the increased weight has a detrimental effect on their gait pattern (and other health issues).

References

- Aisen, M. L., Kerkovich, D., Mast, J., Mulroy, S., Wren, T. Al, Kay, R. M., et al. (2011). Cerebral palsy: clinical care and neurological rehabilitation. *Lancet Neurol.* 10, 844–852. doi:10.1016/S1474-4422(11)70176-4.
- Amiri, P., Hubley-Kozey, C. L., Landry, S. C., Stanish, W. D., and Astephen Wilson, J. L. (2015). Obesity is associated with prolonged activity of the quadriceps and gastrocnemii during gait. *J. Electromyogr. Kinesiol.* 25, 951–8. doi:10.1016/j.jelekin.2015.10.007.
- Bax, M., Goldstein, M., Rosenbaum, P., Leviton, A., Paneth, N., Dan, B., et al. (2005). Proposed definition and classification of cerebral palsy, April 2005. *Dev. Med. Child Neurol.* 47, 571–576. doi:10.1017/S001216220500112X.
- Blakemore, V. J., Fink, P. W., Lark, S. D., and Shultz, S. P. (2013). Mass affects lower extremity muscle activity patterns in children's gait. *Gait Posture* 38, 609–613. doi:10.1016/j.gaitpost.2013.02.002.
- Browning, R. C., and Kram, R. (2007). Effects of obesity on the biomechanics of walking at different speeds. *Med. Sci. Sports Exerc.* 39, 1632–1641. doi:10.1249/mss.0b013e318076b54b.
- Cousins, S. D., Morrison, S. C., and Drechsler, W. I. (2013). Foot loading patterns in normal weight, overweight and obese children aged 7 to 11 years. *J. Foot Ankle Res.* 6, 36. doi:10.1186/1757-1146-6-36.
- Cuthbert, S. C., and Goodheart, G. J. (2007). On the reliability and validity of manual muscle testing: a literature review. *Chiropr. Osteopat.* 15, 4. doi:10.1186/1746-1340-15-4.
- Dallmeijer, A. J., Baker, R., Dodd, K. J., and Taylor, N. F. (2011). Association between isometric muscle strength and gait joint kinetics in adolescents and young adults with cerebral palsy. *Gait Posture* 33, 326–332. doi:10.1016/j.gaitpost.2010.10.092.
- Daniels, L., and Worthingham, C. (1986). *Muscle testing techniques of manual examination*. Philadelphia: WB Saunders.
- Day, S. M., Strauss, D. J., Vachon, P. J., Rosenbloom, L., Shavelle, R. M., and Wu, Y. W. (2007). Growth patterns in a population of children and adolescents with cerebral palsy. *Dev. Med. Child Neurol.* 49, 167–171.
- Eken, M. M., Dallmeijer, A. J., Doorenbosch, C. A., Dekkers, H., Becher, J. G., and Houdijk, H. (2014). Assessment of muscle endurance of the knee extensor muscles in adolescents with spastic cerebral palsy using a submaximal repetitions-to-fatigue protocol. *Arch. Phys. Med. Rehabil.* 95, 1888–94. doi:10.1016/j.apmr.2014.05.010.
- European Union (2014). EU Action Plan on Childhood Obesity 2014-2020. 68. Available at: http://ec.europa.eu/health/nutrition_physical_activity/docs/childhoodobesity_actionplan_

2014_2020_en.pdf [Accessed March 24, 2016].

Finucane, M. M., Stevens, G. A., Cowan, M., Danaei, G., Lin, J. K., Paciorek, C. J., et al. (2011). National, regional, and global trends in body mass index since 1980: Systematic analysis of health examination surveys and epidemiological studies with 960 country-years and 9.1 million participants. *Lancet (London, England)* 377, 557–567. doi:10.1016/S0140-6736(10)62037-5.

Gage, J. R. (2004). “Treatment principles for crouch gait,” in *Treatment of gait problems in cerebral palsy*, ed. J. R. Gage (London: Mac Keith Press), 382–397.

Ghori, G. M., and Luckwill, R. G. (1985). Responses of the lower limb to load carrying in walking man. *Eur. J. Appl. Physiol. Occup. Physiol.* 54, 145–150. doi:10.1007/BF02335921.

Goldberg, S. R., Öunpuu, S., Arnold, A. S., Gage, J. R., and Delp, S. L. (2006). Kinematic and kinetic factors that correlate with improved knee flexion following treatment for stiff-knee gait. *J. Biomech.* 39, 689–698. doi:10.1016/j.jbiomech.2005.01.015.

Gushue, D. L., Houck, J., and Lerner, A. L. (2005). Effects of childhood obesity on three-dimensional knee joint biomechanics during walking. *J. Pediatr. Orthop.* 25, 763–8. doi:00004694-200511000-00012 [pii].

Harding, G. T., Hubley-Kozey, C. L., Dunbar, M. J., Stanish, W. D., and Astephen Wilson, J. L. (2012). Body mass index affects knee joint mechanics during gait differently with and without moderate knee osteoarthritis. *Osteoarthr. Cartil.* 20, 1234–1242. doi:10.1016/j.joca.2012.08.004.

Inman, V. T., Ralston, H. J., and Todd, F. (1981). *Human walking*. Baltimore: Williams & Wilkins.

Kadaba, M. P., Ramakrishnan, H. K., and Wootten, M. E. (1990). Measurement of lower extremity kinematics during level walking. *J. Orthop. Res.* 8, 383–92. doi:10.1002/jor.1100080310.

Kedem, P., and Scher, D. M. (2016). Evaluation and management of crouch gait. *Curr. Opin. Pediatr.* 28, 55–9. doi:10.1097/MOP.0000000000000316.

Lauer, R. T., Johnston, T. E., Smith, B. T., and Lee, S. C. K. (2008). Lower extremity muscle activity during cycling in adolescents with and without cerebral palsy. *Clin. Biomech. (Bristol, Avon)* 23, 442–9. doi:10.1016/j.clinbiomech.2007.11.004.

Liao, H.-F., Jeng, S.-F., Lai, J.-S., Cheng, C.-K., and Hu, M.-H. (1997). The relationship between standing balance and walking function in children with spastic diplegic cerebral palsy. *Dev. Med. Child Neurol.* 39, 106–112.

Mueller, S., Carlsohn, A., Mueller, J., Baur, H., and Mayer, F. (2016). Influence of Obesity on Foot Loading Characteristics in Gait for Children Aged 1 to 12 Years. *PLoS One* 11, e0149924. doi:10.1371/journal.pone.0149924.

Ng, M., Fleming, T., Robinson, M., Thomson, B., Graetz, N., Margono, C., et al. (2014). Global, regional, and national prevalence of overweight and obesity in children and adults during 1980–2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet (London, England)* 384, 766–781. doi:10.1016/S0140-6736(14)60460-8.

Novacheck, T. F., and Gage, J. R. (2007). Orthopedic management of spasticity in cerebral palsy. *Childs. Nerv. Syst.* 23, 1015–1031. doi:10.1007/s00381-007-0378-6.

Novacheck, T., Kroll, G., Gent, G., Rozumalski, A., Beattie, C., and Schwartz, M. (2009). “Orthoses,” in *The identification and treatment of gait problems in cerebral palsy.*, eds. J. Gage, M. Schwartz, S. Koop, and T. Novacheck (Mac Keith Press), 660.

Odding, E., Roebroek, M. E., and Stam, H. J. (2006). The epidemiology of cerebral palsy: incidence, impairments and risk factors. *Disabil. Rehabil.* 28, 183–91. doi:10.1080/09638280500158422.

de Onis, M., Blössner, M., and Borghi, E. (2010). Global prevalence and trends of overweight and obesity among preschool children. *Am. J. Clin. Nutr.* 92, 1257–1264. doi:10.3945/ajcn.2010.29786.1.

Park, E. S., Chang, W. H., Park, J. H., Yoo, J. K., Kim, S. M., and Rha, D. W. (2011). Childhood obesity in ambulatory children and adolescents with spastic cerebral palsy in Korea. *Neuropediatrics* 42, 60–66. doi:10.1055/s-0031-1279724.

Perry, J. (1992). *Gait analysis: Normal and pathological function*. Thorofare, New Jersey: SLACK Incorporated.

Rogers, D. R., and MacIsaac, D. T. (2011). EMG-based muscle fatigue assessment during dynamic contractions using principal component analysis. *J. Electromyogr. Kinesiol.* 21, 811–8. doi:10.1016/j.jelekin.2011.05.002.

Rogozinski, B. M., Davids, J. R., Davis, R. B., Christopher, L. M., Anderson, J. P., Jameson, G. G., et al. (2007). Prevalence of obesity in ambulatory children with cerebral palsy. *J. Bone Joint Surg. Am.* 89, 2421–2426. doi:10.2106/JBJS.F.01080.

Schutte, L. M., Narayanan, U., Stout, J. L., Selber, P., Gage, J. R., and Schwartz, M. H. (2000). An index for quantifying deviations from normal gait. *Gait Posture* 11, 25–31. doi:10.1016/S0966-6362(99)00047-8.

Schwartz, M. H., Viehweger, E., Stout, J., Novacheck, T. F., and Gage, J. R. (2004). Comprehensive treatment of ambulatory children with cerebral palsy: an outcome assessment. *J. Pediatr. Orthop.* 24, 45–53.

Simpson, K. M., Munro, B. J., and Steele, J. R. (2011). Backpack load affects lower limb muscle activity patterns of female hikers during prolonged load carriage. *J. Electromyogr. Kinesiol.* 21, 782–8. doi:10.1016/j.jelekin.2011.05.012.

Taylor, V. H., Forhan, M., Vigod, S. N., McIntyre, R. S., and Morrison, K. M. (2013). The impact of obesity on quality of life. *Best Pract. Res. Clin. Endocrinol. Metab.* 27, 139–146. doi:10.1016/j.beem.2013.04.004.

Trost, J. (2009). “Clinical assessment,” in *The identification and treatment of gait problems in cerebral palsy*, eds. J. Gage, M. Schwartz, S. Koop, and T. Novacheck (London: Mac Keith Press), 181–204.

Visscher, T. L., and Seidell, J. C. (2001). The public health impact of obesity. *Annu. Rev. Public Health* 22, 355–375. doi:10.1146/annurev.publhealth.22.1.355.

Winter, D. A. (1987). *The biomechanics and motor control of human walking*. Waterloo,

520 Canada: University of Waterloo Press.

521 Wolf, S., Loose, T., Schablowski, M., Döderlein, L., Rupp, R., Gerner, H. J., et al. (2006).
522 Automated feature assessment in instrumented gait analysis. *Gait Posture* 23, 331–338.
523 doi:10.1016/j.gaitpost.2005.04.004.

524 World Health Organization (2016a). Report of the commission on ending childhood obesity.
525 68. Available at: <http://www.who.int/end-childhood-obesity/final-report/en/> [Accessed
526 March 24, 2016].

527 World Health Organization (2016b). Available at: <http://www.who.int/> [Accessed March 1,
528 2016].

529

530

In review

Figure legends

Figure 1. Schematic overview of the set-up of the study. The difference scores (walking with added weights vs. weight-free walking) of TD children, all CP children from the CP study group, CP children with adequate strength, and CP children with moderate strength are compared in the current study. In the weighted walking task, 10% of the body weight was added evenly around the waist by means of a black belt. The belt was positioned close to the location of S2 to avoid challenging the equilibrium and ensuring full visibility of the markers of the lower limb Plug-in-Gait marker set.

Figure 2. Mean (+ SE) 3D gait analysis parameters that significantly differed in response to walking with added weight between typically developing children (TD) and children with cerebral palsy (CP) and/or between TD, CP children with adequate strength (aCP) and CP children with moderate strength (mCP). Black bars represent the weight-free trials, white bars represent the weighted walking trials. See Table 2 and 3 for specific significance values. m= meter, % GC= percentage of gait cycle, m/s= meter per second, Gas= gastrocnemius muscle, ampl= amplitude. * $p \leq 0.05$, ** $p \leq 0.01$, *** $p \leq 0.001$

Figure 3. The EMG amplitude difference scores [mean (+ SE)] of the gastrocnemius muscle for the typically developing children (TD) and the children with cerebral palsy (CP) in response to walking with added weights. Comparisons are made to the full CP study group and separately to the two muscle strength CP sub-groups: aCP= adequate lower limb strength, mCP= moderate lower limb strength. Difference scores are expressed as percentage increase/decrease. * $p \leq 0.05$.

554 **Tables**

555 Table 1. Participant characteristics

		CP group		TD group
		Adequate muscle strength group (n=17)	Moderate muscle strength group (n=13)	Typical: typical muscle strength group (n=15)
Average age (SD)		9y 6m ($\pm 1y$ 3m)	6y 5m ($\pm 2y$ 3m)	8y 6m ($\pm 1y$ 3m)
Average weight (SD)		31.8kg (± 10.7 kg)	24.3kg (± 9.3 kg)	29.8kg (± 7.6 kg)
Median (IQR) range of motion (degrees)				
	Hip extension	0° (0° – 0°)	0° (-5° – 0°)	Nr
	Knee extension	0° (0° – 0°)	0° (0° – 5°)	Nr
	Ankle dorsiflexion	0° (0° – 10°)	5° (0° – 10°)	Nr
Selectivity (range 0-2)				
	Knee flexors	2 (2 – 2)	1.5 (1.5 – 2)	Nr
	Knee extensors	2 (2 – 2)	2 (1.5 – 2)	Nr
	Ankle plantarflexors	1.5 (1.25 – 2)	1.5 (1 – 1.5)	Nr
Diagnosis		CP	CP	TD
	Unilateral	9	3	Na
	Bilateral	8	10	Na
GMFCS				
	I	13	4	Na
	II	4	9	Na

556 SD= standard deviation; y= years; m= months; IQR= interquartile range; range of motion= maximal joint angle achieved passively; Nr= normal; CP= cerebral palsy; TD=
557 typically developing; Na= not applicable; GMFCS= Gross Motor Functional Classification Score.

Table 2. Overview of the spatiotemporal, kinematic and kinetic 3DGA parameters that responded significantly different to walking with added weight between typically developing (TD) children and children from the full CP group, CP children with adequate strength (aCP), and CP children with moderate strength (mCP). The values of these parameters for both baseline (weight-free) walking trials and weighted walking trials are presented.

3DGA parameter	TD		CP		aCP		mCP	
	Weight-free (±SE)	Weighted (±SE)	Weight-free (±SE)	Weighted (±SE)	Weight-free (±SE)	Weighted (±SE)	Weight-free (±SE)	Weighted (±SE)
Step length (m)	0.54 (0.02)	0.57 (0.03)	0.46 (0.02)	0.44 (0.02)	0.49 (0.02)	0.48 (0.02)	0.42 (0.02)	0.39 (0.03)
Walking velocity (m/s)	1.17 (0.03)	1.28 (0.06)	1.03 (0.04)	0.94 (0.04)	1.09 (0.04)	1.02 (0.05)	0.94 (0.06)	0.83 (0.06)
Stride length (m)	1.07 (0.03)	1.14 (0.05)	0.95 (0.03)	0.89 (0.04)	1.03 (0.04)	0.96 (0.04)	0.84 (0.04)	0.78 (0.05)
Duration of double support (s)	0.18 (0.01)	0.16 (0.01)	0.18 (0.01)	0.22 (0.01)	0.18 (0.02)	0.21 (0.01)	0.17 (0.02)	0.24 (0.03)
Timing of foot-off (%GC)	59.4 (0.5)	58.9 (0.4)	57.9 (0.7)	60.6 (0.7)	58.48 (0.95)	60.35 (0.89)	57.18 (1.07)	60.96 (1.09)
Pelvic rom, cor (°)	7.3 (0.4)	8.0 (0.4)	11.6 (0.7)	9.2 (0.6)	11.57 (1.05)	9.29 (0.89)	11.52 (0.92)	8.97 (0.88)
Max hip fl swing (°)	35.3 (1.1)	39.1 (1.2)	44.6 (1.6)	44.9 (1.6)	44.15 (2.37)	45.41 (2.24)	45.16 (1.90)	44.28 (2.16)
Hip rom, sag (°)	45.3 (1.1)	50.1 (1.3)	46.9 (1.5)	48.1 (1.8)	47.22 (2.50)	47.26 (2.58)	46.54 (1.37)	49.17 (2.52)
Hip rom, cor (°)	12.6 (0.7)	14.1 (0.6)	14.4 (0.7)	13.6 (0.9)	14.68 (1.12)	13.22 (1.15)	13.98 (0.83)	14.09 (1.34)
Hip rom, trans (°)	21.3 (1.0)	25.6 (1.6)	21.7 (1.1)	20.1 (1.0)	23.47 (1.10)	21.63 (0.85)	19.40 (1.91)	18.03 (1.99)
Max hip fl vel in swing (°/s)	220.5 (6.6)	251.5 (10.0)	223.6 (9.1)	229.8 (11.2)	226.1 (14.2)	229.3 (16.8)	220.5 (10.5)	230.5 (14.6)
Max plfl vel at toe-off (°/s)	161.9 (12.8)	200.9 (10.2)	127.0 (7.8)	121.2 (9.8)	128.4 (10.4)	116.0 (10.5)	125.1 (12.2)	128.1 (18.3)
Max hip ext moment in stance (Nm)	0.50 (0.08)	0.98 (0.07)	0.99 (0.07)	0.80 (0.07)	1.11 (0.09)	0.90 (0.09)	0.79 (0.09)	0.62 (0.10)
Max hip power gen in stance (W)	0.49 (0.09)	0.76 (0.07)	1.36 (0.16)	1.07 (0.14)	1.42 (0.22)	1.18 (0.20)	1.25 (0.21)	0.88 (0.15)
Max hip power abs in stance (W)	-0.69 (0.05)	-0.97 (0.08)	-0.97 (0.09)	-0.94 (0.09)	-0.99 (0.11)	-1.01 (0.12)	-0.92 (0.16)	-0.82 (0.14)
Max hip power gen at toe-off (W)	1.30 (0.09)	1.65 (0.16)	1.15 (0.11)	1.08 (0.14)	1.32 (0.14)	1.33 (0.19)	0.85 (0.16)	0.63 (0.10)
Max knee power gen in stance (W)	0.54 (0.06)	0.99 (0.13)	0.89 (0.14)	0.89 (0.16)	1.00 (0.20)	1.01 (0.23)	0.69 (0.13)	0.67 (0.11)
Max ankle power abs at loading response (W)	-0.63 (0.08)	-0.73 (0.08)	-0.96 (0.09)	-0.82 (0.08)	-0.95 (0.12)	-0.83 (0.10)	-0.97 (0.15)	-0.81 (0.14)

TD= typically developing; CP= cerebral palsy; 3DGA= 3D gait analysis; SE= standard error; rom= range of motion; cor= coronal plane; sag= sagittal plane; trans= transversal plane; max= maximal; fl= flexion; plfl= plantarflexion; ext= extension; vel= velocity; gen= generation; abs= absorption

Table 3. Statistically significant 3DGA parameters and their level of significance per statistical comparison. Comparisons were based on the difference scores between baseline walking and walking with added weights (for the data, see Table 2).

3DGA parameter	Mann-Whitney U values, P-values and Effect Sizes of comparisons based on difference scores (between baseline walking and weighted walking)			
	TD vs. CP	TD vs. aCP	TD vs. mCP	aCP vs. mCP
Step length (m)	89 (.03) / -0.45	ns	ns	ns
Walking velocity (m/s)	81 (.02) / -0.49	ns	ns	ns
Stride length (m)	86 (.02) / -0.47	ns	ns	ns
Duration of double support (s)	74 (.02) / -0.49	ns	ns	ns
Timing of foot-off (% GC)	56 (<.01) / -0.59	40 (<.01) / -0.57	16 (<.01) / -0.68	ns
Pelvic rom, cor (°)	55 (<.01) / -0.61	23 (<.001) / -0.70	32 (<.01) / -0.57	ns
Max hip fl swing (°)	64 (<.01) / -0.58	53 (<.01) / -0.50	11 (.001) / -0.75	ns
Hip rom, sag (°)	113 (.03) / -0.40	ns	ns	ns
Hip rom, cor (°)	108 (.03) / -0.42	ns	ns	ns
Hip rom, trans (°)	92 (.02) / -0.48	ns	ns	ns
Max hip fl vel in swing (°/s)	108 (.03) / -0.42	ns	ns	ns
Max plfl vel at toe-off (°/s)	91 (.02) / -0.48	ns	ns	ns
Max hip ext moment in stance (Nm)	28 (<.001) / -0.70	15 (<.001) / -0.75	13 (<.01) / -0.66	ns
Max hip power gen in stance (W)	82 (.02) / -0.47	ns	ns	ns
Max hip power abs in stance (W)	89 (.03) / -0.44	ns	ns	ns
Max hip power gen at toe-off (W)	88 (.03) / -0.44	ns	ns	ns
Max knee power gen in stance (W)	89 (.03) / -0.40	ns	ns	ns
Max ankle power abs at loading response (W)	96 (.04) / -0.42	ns	ns	ns
Mean EMG ampl gastroc increase/decrease (%)	108 (.03) / -0.42	ns	ns	ns

Data is presented as follows; U-value (p-value) / Effect size. 3DGA= 3D gait analysis; TD= typically developing; CP= cerebral palsy study group; aCP= adequate strength CP group; mCP= moderate strength CP group; ns= not significant; na= not applicable; rom= range of motion; cor= coronal plane; sag= sagittal plane; trans= transversal plane; max= maximal; fl= flexion; plfl= plantarflexion; ext= extension; vel= velocity; gen= generation; abs= absorption; ampl= amplitude; gastroc= gastrocnemius muscle

Figure 1.TIF




Figure 2.TIF




Figure 3.TIF

